

## Colon Surgery

Patients undergo colon surgery for a number of conditions including: colorectal cancer, polyps, inflammatory bowel disease (Crohn's disease and ulcerative colitis), colonic inertia, stricture (narrowing) of the colon and diverticular disease; surgery to remove all or part of your colon is known as colectomy.

Traditional "open" colon surgery procedures may require a single long abdominal incision. This usually results in an average hospital stay of a week or more and 6 weeks of recovery. Less invasive options are available to many patients facing colon surgery. The most common of these is laparoscopic surgery, in which smaller incisions are used.

## The Colon

The colon refers to the large intestine, which is the lower part of the digestive tract. The intestine is a long, tubular organ consisting of the small intestine, the colon (large intestine) and the rectum, which is the last part of the large bowel. After food is swallowed, it begins to be digested in the stomach and then empties into the small intestine, where the nutritional part of the food is absorbed. The remaining waste moves through the colon to the rectum and is expelled from the body. The colon and rectum absorb water and hold the waste until it is ready to be expelled through a process known as defaecation.

## Laparoscopic Right Hemicolectomy

A technique known as minimally invasive laparoscopic colon surgery allows surgeons to perform many common colon procedures through small incisions. Depending on the type of procedure, patients may leave the hospital in a few days and return to normal activities more quickly than patients recovering from open surgery.

In laparoscopic right hemicolectomy, surgeons operate through 3 or 4 small openings (each about a quarter inch), while watching an enlarged image of the patient's internal organs on a television monitor. In some cases, one of the small openings may be lengthened to 2 or 3 inches to extract the specimen and complete the procedure.

The term right colectomy refers to the removal of the right (more proximal) side of the colon; typically about 1/3 of the colon is removed. The remaining large bowel is normally sufficient to ensure that no significant changes to the bowel function (frequency, consistency and pattern) occur after surgery.

## Advantages Of Laparoscopic Colon Resections

Results may vary depending upon the type of procedure and patient's overall condition.

Common advantages are:

- Less postoperative pain
- Generally shorter hospital stay
- Faster return to solid-food diet
- May result in a quicker return of bowel function
- Quicker return to normal activity

Enquiries: 07500870587 or 01519295181

[enquiries@wirralsurgeon.co.uk](mailto:enquiries@wirralsurgeon.co.uk)

[www.wirralsurgeon.co.uk](http://www.wirralsurgeon.co.uk)

- Improved cosmetic results
- Less chance for developing an incisional hernia or adhesions
- Lower rates of wound infection

### **Are You A Candidate For Laparoscopic Colon Resection?**

Although laparoscopic colon resection has many benefits, it may not be appropriate for some patients. Typically, laparoscopic surgery might not be suitable for patients with:

- Significant obesity (BMI over 35)
- A history of prior abdominal surgery causing dense scar tissue (adhesions)
- Large tumors
- Certain respiratory or cardiac conditions
- Previous organ transplantation
- Disorders of coagulation (blood clotting)

Each patient will be offered a thorough and objective evaluation during her/his initial consultation with us to find out if the technique is appropriate for her/his condition.

### **Diagnostic tests**

Most diseases of the colon are diagnosed with one of two tests: a colonoscopy or CT virtual colonoscopy. A colonoscope is a soft, bendable tube about the thickness of the index finger, which is inserted into the anus (back passage) and then advanced through the entire large intestine (see separate information sheet on Colonoscopy). A CT virtual colonoscopy is a special radiological investigation (scan) where the entire large intestine is imaged by the use of a CT scan and intravenous and intraluminal (within the bowel) contrast media. In addition, a CT scan of the chest, abdomen and pelvis may be necessary in cases of bowel cancer, inflammatory bowel disease or diverticular disease.

### **Preparation For Surgery**

Pre-operative preparation includes blood tests, medical evaluation, chest x-ray and an electrocardiogram (ECG), depending on your age and medical condition. In addition, patients with significant co-morbidity might be asked to undergo lung function tests and an echocardiogram (ultrasound examination of the heart). In high-risk patients a formal anaesthetic evaluation and cardio-pulmonary exercise (CPEX) testing will be performed. Blood transfusion and/or blood products may be needed depending on your condition and the amount of blood loss during surgery.

It is recommended that you shower the night before or morning of the operation.

In general, no formal bowel preparation (emptying) is required for a right hemicolectomy. In special cases, we might want your colon and rectum to be completely empty before surgery. If this is the case, you must drink a special cleansing solution. You may be on several days of clear liquids, laxatives and enemas prior to the operation. If you are unable to take the preparation please let us know in advance. If you do not complete the preparation, it may be unsafe to undergo the surgery and it may have to be rescheduled.

After midnight the night before the operation, you should not eat or drink anything except

Enquiries: 07500870587 or 01519295181

[enquiries@wirralsurgeon.co.uk](mailto:enquiries@wirralsurgeon.co.uk)

[www.wirralsurgeon.co.uk](http://www.wirralsurgeon.co.uk)

medications that your surgeon has told you are permissible to take with a sip of water the morning of surgery.

Drugs such as aspirin, blood thinners, anti-inflammatory medications (arthritis medications) and Vitamin E will need to be stopped temporarily for several days to a week prior to surgery. Diet medication or St. John's Wort should not be used for the two weeks prior to surgery.

The amount of alcohol you drink can affect you during and after your surgery. It is important that you talk with us about your alcohol intake so that we can plan your care. Stopping alcohol suddenly can cause seizures, delirium, and death. If we know you are at risk for these complications, we can prescribe medications to help prevent them. If you drink alcohol regularly, you may be at risk for other complications during and after surgery. These include bleeding, infections, heart problems, and a longer hospital stay. Here are things you can do to prevent problems before your surgery:

- Be honest with us about how much alcohol you drink
- Try to stop drinking alcohol once your surgery is planned. If you develop a headache, nausea, increased anxiety, or cannot sleep after you stop drinking, tell your doctor right away. These are early signs of alcohol withdrawal and can be treated.
- Tell us if you cannot stop drinking.

People who smoke can have breathing problems when they have surgery. Stopping even for a few days before surgery can help. Please quit smoking for at least 48 hours before surgery and arrange for any help you may need at home.

### **How Is Laparoscopic Colon Resection Performed?**

"Laparoscopic" or "Key-hole" surgery describes the techniques a surgeon uses to gain access to the abdominal cavity. A specialized camera called a laparoscope (a tiny telescope connected to a video camera) is inserted through a port (a narrow hollow tube like instrument) placed through the abdominal wall. At the beginning of the procedure, the abdomen is inflated with carbon dioxide gas to provide a working space for the surgeon. The laparoscope transmits images inside the abdominal cavity, giving the surgeon a magnified view of the patient's internal organs on a television monitor.

Several other ports are subsequently inserted to allow the surgeon to work inside and remove part of the colon. At the end of the procedure the resected bowel is typically removed by lengthening one of the small port incisions, most commonly the one just below the umbilicus (belly button).

### **What Happens If The Operation Cannot Be Performed Or Completed By The Laparoscopic Method?**

In a number of patients the laparoscopic method cannot be performed. Factors that may increase the possibility of choosing or converting to the "open" procedure may include:

- Obesity
- A history of prior abdominal surgery causing dense scar tissue
- Inability to visualize organs
- Bleeding problems during the operation

Enquiries: 07500870587 or 01519295181

[enquiries@wirralsurgeon.co.uk](mailto:enquiries@wirralsurgeon.co.uk)

[www.wirralsurgeon.co.uk](http://www.wirralsurgeon.co.uk)

- Large tumors

The decision to perform the open procedure is a judgment decision made by your surgeon either before or during the actual operation. When the surgeon feels that it is safest to convert the laparoscopic procedure to an open one; this is not a complication, but rather sound surgical judgment. The decision to convert to an open procedure is strictly based on patient safety.

### **What Should I Expect After The Surgery?**

After the operation, it is important to follow our instructions. Although many people feel better in a few days, remember that your body needs time to heal.

When you wake up after your surgery, you will be in the Recovery Area. You will stay there until you are awake and your pain is under control. Most people return to their ward after 2-3 hours. Patients with significant pre-existing medical conditions will be transferred to the High Dependency Unit (HDU) or Intensive Care Unit (ICU) instead.

You will receive oxygen through a thin tube called a nasal cannula that rests below your nose. A nurse will be monitoring your body temperature, pulse, blood pressure, and oxygen levels.

You will have a Foley® catheter in your bladder to monitor the amount of urine you are making. You will also have compression stockings on your lower legs to help your circulation. They will be taken off when you are able to walk.

You will be given medications to control your pain and keep you comfortable. There are different ways that these medications can be given:

- Epidural catheter: some people may get pain medication through an epidural catheter in their spine
- Nerve block: some patients may get a nerve block before or during surgery. In a nerve block, your doctor injects medication into some of your nerves to reduce pain after surgery
- Intravenous (IV) medications: some people may get pain medication straight into a vein through their IV line
- Oral medications: some patients may get oral pain medications (medication that's swallowed, such as pills).

You may have one or more of these after your surgery. They're all effective methods to control your pain, and the Anaesthetist will talk with you before choosing the best one(s) for you.

You will be offered fluids shortly after your surgery. Following this you should then be able to eat and drink normally within 1-3 days.

Your bowel will usually start to make sounds after 1 or 2 days and you may have a bowel movement after 3 to 4 days. However, if this does not happen you should not be too worried. Bowel movements are different from one person to another. Usually, the first sign that your bowel is beginning to work is when you pass wind. You may then find that your bowel habits become unpredictable, with a sense of urgency. The colour and consistency of your bowel motions may also vary. Your bowel habit should settle down, and your bowel

Enquiries: 07500870587 or 01519295181

[enquiries@wirralsurgeon.co.uk](mailto:enquiries@wirralsurgeon.co.uk)

[www.wirralsurgeon.co.uk](http://www.wirralsurgeon.co.uk)

control will improve with time.

We only use absorbable wound stitches so no stitches need to be removed after operation; this also ensures a very good cosmetic appearance. The wound becomes waterproof 24 hours after surgery, so you can shower or bath without fear after the second post-operative day.

You are encouraged to be out of bed the day after surgery and to walk. This will help diminish the soreness in your muscles. You will probably be able to get back to most of your normal activities in one to two weeks' time. These activities include driving, walking up the stairs, working and engaging in sexual intercourse.

In most cases a follow-up appointment is organized within 2 weeks after your operation.

### **What Complications Can Occur?**

Complications after colonic surgery are unfortunately not uncommon and include:

- Bleeding
- Infection (of the wounds, inside the abdomen, chest, bladder)
- A leak where the colon was connected back together (anastomotic leak or dehiscence); this is one of the most serious complications after colon surgery. Although some cases can be treated with antibiotics, bowel rest and/or drainage of abscess via a small tube inserted under radiologic control, most anastomotic leaks necessitate return to theatre and formation of a stoma (an opening on the surface of the abdomen which has been created surgically to divert the flow of faeces)
- Injury to adjacent organs such as the small intestine, ureter, or bladder
- Blood clots in deep veins in your legs (deep vein thrombosis) that may travel to your lungs (pulmonary embolism)
- Incisional hernia (smaller rate than with conventional surgery)
- Adhesions (much smaller risk than with open surgery).

It is important for you to recognize the early signs of possible complications. Contact us on 07968228831 or present to the Accident and Emergency Department if you notice severe abdominal pain, fevers, chills, or rectal bleeding.

### **When to Call Your Surgeon**

Be sure to call us if you develop any of the following:

- Persistent fever over 38°C
- Bleeding from the rectum
- Increasing abdominal swelling
- Pain that is not relieved by your medications
- Persistent nausea and/or vomiting
- Chills
- Persistent cough or shortness of breath
- Purulent drainage (pus) from any incision
- Redness surrounding any of your incisions that is worsening or getting bigger
- You are unable to eat or drink liquids.

Enquiries: 07500870587 or 01519295181

[enquiries@wirralsurgeon.co.uk](mailto:enquiries@wirralsurgeon.co.uk)

[www.wirralsurgeon.co.uk](http://www.wirralsurgeon.co.uk)